

AD-A160 612

ANALYSES OF EXTRATERRESTRIAL MATERIALS IN TERRESTRIAL
SEDIMENTS(U) CALIFORNIA UNIV LOS ANGELES DEPT OF
GEOLOGY F T KYTE 1983 N00014-81-K-0688

1/1

UNCLASSIFIED


F/G 3/2

NL

END

FILMED

DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A160 612

ABSTRACT OF THE DISSERTATION

Analyses of Extraterrestrial Materials

in Terrestrial Sediments

1983


by

Frank T. Kyte

Doctor of Philosophy in Geology

University of California, Los Angeles, 1983

Professor John T. Wasson, Chair



Analyses of Cretaceous-Tertiary boundary sediments from Denmark and 2 Pacific sites (465A and GPC-3) have aided in demonstrating that the high concentrations of Ir in this boundary is a global phenomenon. This is consistent with the occurrence of a major accretionary event at this time, one which may have resulted in the massive terminal Cretaceous extinctions. The presence of high Ir concentrations in a mid-Pacific carbonate sequence (465A) and abyssal clay (GPC-3) indicates that this phenomenon is not restricted to shallow continental shelf regions, but occurs in a variety of environments. Relative abundances of the siderophiles Os, Re, Ir, Pt, Ni, Co, Fe, Pd, and Au in the Danish samples are similar to chondritic meteoritic abundances (within a factor of 3, relative to Ir) and indicate an extraterrestrial source. However, variations in siderophile abundances between sites and even between samples at each site indicate significant fractionation either during accretion, depo-

DTIC FILE COPY

DISTRIBUTION STATEMENT A

Approved for public release
Distribution Unlimited

x

85 09 11 031

sition, or diagenetic alteration. These fractionations make it impossible at present to use siderophiles to establish a link between the accreting object and a known meteorite group. The best evidence for the impact of a compact asteroidal object (as opposed to accretion of fragmented materials) would be characterization of an ejecta component in boundary clays, but conclusive evidence is still lacking.

Anomalous concentrations of Ir and Au in a Late Pliocene (~2.3 Ma) section from the Antarctic Basin was the first such discovery since the KT boundary. Associated coarse-grained debris contains a vesicular impact melt and unmelted clasts of the projectile. Chemical and petrographic evidence indicates that this horizon resulted from the oceanic impact of an asteroid similar to mesosiderite or howardite meteorites. Estimates of the projectile diameter are in the range of 100 to 500 m.

Analyses of cosmic spherules separated from abyssal clays indicate that most stony spherules are destroyed by weathering before burial to a depth of 0.5 m. At least half of the iron spherules may be derived from chondritic precursors; these contain a majority of the total siderophiles. Revised influx rates suggest that >10% of the materials in the size range 10^{-5} to 10^6 g survive atmospheric entry without vaporization.

SEARCHED		INDEXED	
SERIALIZED		FILED	
APR 1971			
FBI - NEW YORK			
Dist			
A-1			

Acknowledgements

Chapter 2 originally appeared in Nature 288, 651-656 (1980), by Frank T. Kyte, Zhiming Zhou, and John T. Wasson. Chapter 4 originally appeared in Nature 292, 417-420 (1981), by Frank T. Kyte, Zhiming Zhou, and John T. Wasson. Chapter 6 originally appeared in Geol. Soc. America Spec. Paper 190, 235-242 (1982), by Frank T. Kyte and John T. Wasson.

Special thanks belong to J. T. Wasson whose supervision and direction has greatly influenced this research project. Special thanks also to M. B. Blanchard who first introduced me to analysis of extraterrestrial materials and D. E. Brownlee whose numerous conversations, criticisms, and collaboration has greatly increased my understanding of extraterrestrial particles.

Many thanks go to Zhou Z. for his collaboration during the preliminary phases of this research, particularly for his tireless efforts in developing a radiochemical procedure for analyses of noble metals. Many thanks also to G. W. Kallemeyn for his essential assistance in neutron activation analysis, and to J. N. Grossman for the numerous computer programs which have been invaluable to our entire research group. Their assistance has been an essential part of this research project.

I also wish to thank R. S. Wong, P. Tilke, D. J. Malvin, P. S. Rosener, D. N. Shirley, J. Willis, K. L. Rasmussen, and

Zhang Z.-X., my fellow graduate students, post-docs, scholars and lab helpers, for their assistance and for the many helpful discussions, criticisms and comments they provided.

Other thanks go to G. Chen, G. No, J. Pai, P. L. Tu, and A. Young for help in the preparation of manuscripts, to R. Jones for help in learning to use the electron microprobe, and to J. L. Ghafourpour, V. Doyle-Jones and J. Sells for the skillful drafting of many figures.

The samples from Stevns Klint, Denmark were kindly provided by A. R. Loeblich and H. Tappan. The samples from DSDP Hole 465A were supplied with the assistance of W. R. Riedel and L. Garifal. Samples of GPC-3 were supplied by Woods Hole Oceanographic Institute with the assistance of B. H. Corliss, J. Broda and B. Wooding. The samples from E13-3 and E13-4 were supplied by Florida State University with very helpful assistance from D. S. Cassidy and F. A. Kaharoeddin.

This research was mainly supported by the Office of Naval Research contract N00014-81-K-0688 and the National Science Foundation contract OCE 82-08197.

END

FILMED

12-85

DTIC